

Districts.	Accumulated departures.		Districts.	Accumulated departures.	
	Total.	Average.		Total.	Average.
West Gulf	+ 2.8	+ 0.7	New England	- 3.7	- 0.9
Ohio Valley and Tenn.	+ 2.0	+ 0.5	Middle Atlantic	- 2.0	- 0.5
Lower Lake	+ 2.0	+ 0.5	South Atlantic	- 2.2	- 0.6
Upper Lake	+ 9.8	+ 2.4	Florida Peninsula	- 9.7	- 2.4
North Dakota	+ 6.0	+ 1.5	East Gulf	- 4.8	- 1.2
Upper Mississippi	+ 12.7	+ 3.2			
Missouri Valley	+ 14.6	+ 3.6			
Northern Slope	+ 8.9	+ 2.2			
Middle Slope	+ 16.5	+ 4.1			
Abilene (southern Slope) ..	+ 10.0	+ 2.5			
Southern Plateau	+ 3.8	+ 1.0			
Middle Plateau	+ 4.9	+ 1.2			
Northern Plateau	+ 17.4	+ 4.4			
North Pacific	+ 0.9	+ 0.2			
Middle Pacific	+ 2.2	+ 0.6			
Southern Pacific	+ 4.5	+ 1.1			

The limit of freezing weather is shown on Chart VI by the isotherm of minimum 32°, and the limit of frost by the isotherm of minimum 40°.

MOISTURE.

The quantity of moisture in the atmosphere at any time may be expressed by the weight of the vapor coexisting with the air contained in a cubic foot of space, or by the tension or pressure of the vapor, or by the temperature of the dew-point. The mean dew-points for each station of the Weather Bureau, as deduced from observations made at 8 a. m. and 8 p. m., daily, are given in Table I.

The rate of evaporation from a special surface of water on muslin at any moment determines the temperature of the wet-bulb thermometer, but a properly constructed evaporimeter may be made to give the quantity of water evaporated from a similar surface during any interval of time. Such an evaporimeter, therefore, would sum up or integrate the effects of those influences that determine the temperature as given by the wet bulb; from this quantity the average humidity of the air during any given interval of time may be deduced.

Measurements of evaporation within the thermometer shelters are difficult to make so as to be comparable at temperatures above and below freezing, and may be replaced by computations based on the wet-bulb temperatures. The absolute amount of evaporation from natural surfaces not protected from wind, rain, sunshine, and radiation, are being made at a few experimental stations and will be discussed in special contributions.

Sensible temperatures.—The sensation of temperature experienced by the human body and ordinarily attributed to the condition of the atmosphere depends not merely on the temperature of the air, but also on its dryness, on the velocity of the wind, and on the suddenness of atmospheric changes, all combined with the physiological condition of the observer. A complete expression for the relation between atmospheric conditions and nervous sensations has not yet been obtained.

PRECIPITATION.

[In inches and hundredths.]

The distribution of precipitation for the current month, as determined by reports from about 2,500 stations, is exhibited on Chart III. The numerical details are given in Tables I, II, and III. The total precipitation for the current month was heaviest (from 6 to 18 inches) in the coastal and mountainous regions of northern California, Washington, and Oregon. Areas of 8 to 10 inches were reported from eastern Nebraska, inclosed within a much larger region of over 6 inches.

The larger values at regular stations were: Eureka, 11.1; Astoria, 9.2; Neahbay, 7.9; Greenbay, 7.5.

The diurnal variation, as shown by tables of hourly means

of the total precipitation, deduced from self-registering gauges kept at the regular stations of the Weather Bureau, is not now tabulated.

The current departures from the normal precipitation are given in Table I, which shows that precipitation was in excess over the upper Lake Region, upper Mississippi, and lower Missouri valleys, Manitoba, Washington, Oregon, and northern California. It was deficient in the Atlantic States. The large excesses were: Eureka, 7.0; Dubuque, 5.0; Winnipeg, 4.3; Dodge City, 3.9; La Crosse, 3.6; Greenbay and Astoria, 3.3. The large deficits were: Fort Smith, 4.6; Louisville, 4.2; Vicksburg, 3.6; Palestine, 3.4; Atlanta, 3.1; Nantucket and Hatteras, 3.0.

The average departure for each district is also given in Table I. By dividing these by the respective normals the following corresponding percentages are obtained (precipitation is in excess when the percentages of the normals exceed 100):

Above the normal: Lower Lake, 109; upper Lake, 147; North Dakota, 160; upper Mississippi, 130; Missouri Valley, 138; northern Slope, 106; middle Slope, 109; middle Plateau, 145; northern Plateau, 120; north Pacific, 127; middle Pacific, 236.

Normal: Southern Plateau, 100.

Below the normal: New England, 38; middle Atlantic, 37; south Atlantic, 42; Florida Peninsula, 46; east Gulf, 61; west Gulf, 56; Ohio Valley and Tennessee, 57; southern Plateau, (Abilene), 41; southern Pacific, 92.

The years of greatest and least precipitation for April are given in the REVIEW for April, 1890. The precipitation for the current month was the greatest on record at: Williston, 2.86; St. Paul, 5.63; Greenbay, 5.48; La Crosse, 5.84; Minneapolis, 5.12; Huron, 6.17; Sioux City, 6.16; Dubuque, 7.80; Topeka, 4.00; Dodge City, 5.50; Winnemucca, 1.95; Eureka, 11.13; Point Reyes Light, 4.20; Fresno, 2.82. It was the least on record at: Eastport, 0.86; New Haven, 1.19; Nantucket, 0.62; Narragansett Pier, 1.38; Woods Hole, 1.33; Harrisburg, 1.19; Washington, 1.07; Cincinnati, 0.59; Indianapolis, 1.27; Louisville, 0.40; Lexington, 0.40; Wilmington, 0.64; Atlanta, 0.58; Fort Smith, 0.46.

The total accumulated monthly departures from normal precipitation from January 1 to the end of the current month are given in the second column of the following table; the third column gives the ratio of the current accumulated precipitation to its normal value.

Districts.	Accumulated departures.	Accumulated precipitation.	Districts.	Accumulated departures.	Accumulated precipitation.
Lower Lakes	+ 1.00	109	New England	- 2.80	83
North Dakota	+ 2.30	157	Middle Atlantic	- 1.80	85
Northern Slope	+ 0.50	114	South Atlantic	- 3.70	77
Middle Plateau	+ 0.90	116	Florida Peninsula	- 0.60	94
North Pacific	+ 4.40	120	East Gulf	- 4.00	80
Middle Pacific	+ 2.40	114	West Gulf	- 2.50	83
			Ohio Valley and Tenn.	- 5.60	68
			Upper Lakes	- 0.70	92
			Upper Mississippi	- 1.30	86
			Missouri Valley	- 0.50	93
			Middle Slope	- 1.50	73
			Abilene (southern Slope) ..	- 2.90	54
			Southern Plateau	- 0.40	79
			Northern Plateau	- 1.10	85
			South Pacific	- 1.00	79

Details as to excessive precipitation are given in Tables XII and XIII.

The total monthly snowfall at each station is given in Table II. Its geographical distribution is shown on Chart VI. The southern limit of freezing temperatures and possible snow is shown on this chart by the isotherm of minimum 32°. The isotherm of minimum 40°, namely, the air temperature within the thermometer shelter, is also given on this chart,